



Contribution ID: 157

Type: **oral**

Imaging The Renner-Teller Effect Using Laser-Induced Electron Diffraction

Thursday 4 July 2019 16:45 (15 minutes)

We employ laser-induced electron diffraction (LIED) coupled with state-of-the-art quantum dynamical calculations to reveal a linear-to-bent transition that is enabled by the Renner-Teller effect in field-dressed molecules investigated with a single-pulse pump-probe excitation and measurement.

Authors: Dr AMINI, Kasra (ICFO); Dr STEINLE, Tobias (ICFO); Dr SCLAFANI, Michele (ICFO); Mr SANCHEZ, Aurelien (ICFO); Mr LIU, Xinyao (ICFO); Prof. LE, A.T. (Kansas State University); Dr STEINMETZER, Johannes (Friedrich-Schiller University); Prof. PFEIFFER, Thomas (Max-Planck-Institut für Kernphysik); Prof. MOSHAMMER, Robert (Max-Planck-Institut für Kernphysik); Prof. ULLRICH, Joachim (Max-Planck-Institut für Kernphysik); Prof. LEWENSTEIN, Maciej (ICFO); Prof. MOSZYNSKI, Robert (University of Warsaw); Prof. DE ABAJO, Javier (ICFO); Prof. LIN, C.D. (Kansas State University); Prof. GRÄFE, Stefanie (Friedrich-Schiller University); Prof. BIEGERT, Jens (ICFO)

Presenter: Dr AMINI, Kasra (ICFO)

Session Classification: New approaches for measuring atomic and molecular dynamics